

Inseparability criteria for bipartite quantum states

E. Shchukin* and W. Vogel†

Arbeitsgruppe Quantenoptik, Institut für Physik, Universität Rostock, D-18051 Rostock, Germany

We provide necessary and sufficient conditions for the partial transposition of bipartite harmonic quantum states to be nonnegative. The conditions are formulated as an infinite series of inequalities for the moments of the state under study. The violation of any inequality of this series is a sufficient condition for entanglement. Previously known entanglement conditions are shown to be special cases of our approach.

PACS numbers: 03.67.Mn, 03.65.Ud, 42.50.Dv

Entanglement plays a key role in the rapidly developing field of quantum information processing. In this context it is important to provide methods for characterizing entangled quantum states on the basis of observable quantities. However, already in seemingly simple cases this problem turns out to be rather complex. Even for a two-party harmonic-oscillator system so far there exists no complete characterization of entanglement to be used in experiments.

For characterizing entanglement, that is inseparability, of the density operator of a bipartite continuous variable system, one may use the Peres-Horodecki condition [1, 2, 3]. A sufficient condition for entanglement consists in the negativity of the partial transposition (NPT) of the quantum state of the two-party system. To characterize NPT for such a system completely, however, to our best knowledge is still an unsolved problem.

A sufficient condition for the NPT has been proposed by Simon [4]. It is based on second-order moments of position and momentum operators. For the special case of Gaussian states the resulting entanglement criterion has been shown to be necessary and sufficient. Another inseparability condition based on second moments has been derived without explicitly using the NPT condition [5]. This condition is also complete for the characterization of entanglement of two-mode Gaussian states. The latter approach has been extended to special higher-order moments [6, 7] and even to more general operator functions [8].

The complexity of the problem under study may become clear when we go back to a related but simpler problem. It consists in the characterization of nonclassical effects based on the negativity of the Glauber-Sudarshan P -function. Only recently the problem was solved of how to characterize the nonclassicality in terms of observable quantities. This requires an infinite hierarchy of conditions formulated either in terms of characteristic functions [9] or in terms of observable moments [10, 11].

In the present contribution we will further develop the concept of the complete characterization of single-mode nonclassicality with the aim to characterize the entanglement of bipartite continuous variable quantum states. Based on the NPT condition we derive a hierarchy of

necessary and sufficient conditions for the NPT in terms of observable moments. Even though this only leads to sufficient conditions for entanglement, it can be applied to a variety of quantum states. It does not only contain as a special case the condition of Simon [4], but also other types of inequalities [5, 6, 7, 8]. For the sake of strictness of the conclusions we restrict our attention to a two-mode harmonic oscillator. Hence the derived conditions are useful for radiation modes, motional states of trapped ions and related systems. The further extension of our approach to more general continuous variable systems seems feasible, but this requires some further investigations.

A bipartite quantum state $\hat{\rho}$ is called separable if it is a convex combination of factorizable states,

$$\hat{\rho} = \sum_{n=0}^{+\infty} p_n \hat{\rho}_1^{(n)} \otimes \hat{\rho}_2^{(n)}, \quad (1)$$

where $\hat{\rho}_1^{(n)}$ and $\hat{\rho}_2^{(n)}$ are states of the first and the second part respectively. The p_n are nonnegative numbers satisfying $\sum_{n=0}^{+\infty} p_n = 1$ and the series in (1) is convergent in the trace norm. A state which cannot be represented in the form (1) is called entangled.

It is well known that the full transposition of any (one-partite or multi-partite) state is again a quantum state. But in the multipartite case, along with the notion of the full transposition, there is a notion of partial transposition, which transposes only some parts of the state and leaves the others unchanged. In contrast to the full transposition, partial transposition does not necessarily map a quantum state onto a quantum state. Clearly, the partially transposed density operator $\hat{\rho}^{\text{PT}}$ of a separable state $\hat{\rho}$, Eq. (1), is again a quantum state (we consider transposition of the second part; our results are unchanged if we transpose the first part):

$$\hat{\rho}^{\text{PT}} = \sum_{n=0}^{+\infty} p_n \hat{\rho}_1^{(n)} \otimes \hat{\rho}_2^{(n)\text{T}}, \quad (2)$$

since $\hat{\rho}_2^{(n)\text{T}}$ is a quantum state for all n and so is $\hat{\rho}^{\text{PT}}$ according to the expansion (2). This property of separable states allows us to use the NPT as a sufficient condition for inseparability. This condition is referred to as

Peres-Horodecki condition [1, 2, 3]. In general the partial transposition of a density operator may no longer be nonnegative defined. It is this property that we will use for the derivation of our hierarchy of conditions.

Remember that a Hermitian operator $\hat{A}$ is called non-negative if

$$\langle \psi | \hat{A} | \psi \rangle \geq 0 \quad (3)$$

holds for all states $|\psi\rangle$. The left hand side of this inequality can be written as

$$\langle \psi | \hat{A} | \psi \rangle = \text{Tr}(\hat{A} |\psi\rangle \langle \psi|). \quad (4)$$

The nonnegative operator $|\psi\rangle \langle \psi|$ can be represented in the form $\hat{f}^\dagger \hat{f}$, where $\hat{f}$ may be written as

$$\hat{f} = |\text{vac}\rangle \langle \psi|. \quad (5)$$

The state $|\text{vac}\rangle$ is the vacuum state, in our bipartite case it reads, in the Fock basis, as $|\text{vac}\rangle = |00\rangle$. Any pure bipartite state $|\psi\rangle$ can be expressed as

$$|\psi\rangle = \hat{g}^\dagger |00\rangle, \quad (6)$$

for an appropriate operator function $\hat{g} = g(\hat{a}, \hat{b})$, where $\hat{a}$ and $\hat{b}$ are the annihilation operators of the first and the second mode respectively. Hence the operator (5) is of the form

$$\hat{f} = |00\rangle \langle 00| \hat{g}. \quad (7)$$

The two-mode vacuum density operator may be expressed in the normally-ordered form,

$$|00\rangle \langle 00| \equiv : \exp(-\hat{a}^\dagger \hat{a} - \hat{b}^\dagger \hat{b}) :, \quad (8)$$

with $:\dots:$ denoting the normally-ordering prescription. By combining Eqs (7) and (8) it immediately follows that the normally-ordered form of the resulting operator $\hat{f}$ exists. Hence we conclude: *a Hermitian operator is nonnegative if and only if for any operator $\hat{f}$ whose normally-ordered form exists the inequality*

$$\langle \hat{f}^\dagger \hat{f} \rangle_{\hat{A}} = \text{Tr}(\hat{A} \hat{f}^\dagger \hat{f}) \geq 0 \quad (9)$$

is satisfied.

Now we can use the inequality (9) to apply the Peres-Horodecki condition, which gives the following result. *For any separable state $\hat{\varrho}$ the inequality*

$$\langle \hat{f}^\dagger \hat{f} \rangle^{\text{PT}} \equiv \langle \hat{f}^\dagger \hat{f} \rangle_{\hat{\varrho}^{\text{PT}}} = \text{Tr}(\hat{\varrho}^{\text{PT}} \hat{f}^\dagger \hat{f}) \geq 0 \quad (10)$$

holds true for any operator $\hat{f}$ whose normally-ordered form exists. According to the latter assumption the operator $\hat{f}$ can be expanded as

$$\hat{f} = \sum_{n,m,k,l=0}^{+\infty} c_{nmkl} \hat{a}^{\dagger n} \hat{a}^m \hat{b}^{\dagger k} \hat{b}^l. \quad (11)$$

Upon substituting this expansion into the inequality (10) we get the condition

$$\langle \hat{f}^\dagger \hat{f} \rangle^{\text{PT}} = \sum_{\substack{n,m,k,l=0 \\ p,q,r,s=0}}^{+\infty} c_{pqrs}^* c_{nmkl} M_{pqrs,nmkl} \geq 0, \quad (12)$$

for the moments of the partial transposition

$$M_{pqrs,nmkl} = \langle \hat{a}^{\dagger q} \hat{a}^p \hat{a}^{\dagger n} \hat{a}^m \hat{b}^{\dagger s} \hat{b}^r \hat{b}^{\dagger k} \hat{b}^l \rangle^{\text{PT}}. \quad (13)$$

The left hand side of the inequality (12) is a quadratic form with respect to the coefficients c_{nmkl} of the expansion (11). The Sylvester criterion states that the inequality (12) holds for all c_{nmkl} if and only if all the main minors of the from (12) are nonnegative. To write these minors explicitly we need to find the relation between the moments (13) of the partially transposed state and those of the original one.

It is convenient to number multi-indices (n, m, k, l) by a single number so that the moments $M_{pqrs,nmkl}$ be numbered by two indices like matrix elements are. The problem of numbering the multi-indices (n, m, k, l) is equivalent to the ordering of the set $\{(n, m, k, l)\}$ of all such multi-indices. Then we can use the ordinal number of a multi-index (n, m, k, l) in the ordered sequence of multi-indices. We use the following order: for any two multi-indices $\mathbf{u} = (n, m, k, l)$ and $\mathbf{v} = (p, q, r, s)$

$$\mathbf{u} < \mathbf{v} \Leftrightarrow \begin{cases} |\mathbf{u}| < |\mathbf{v}|, \text{ or} \\ |\mathbf{u}| = |\mathbf{v}| \text{ and } \mathbf{u} <' \mathbf{v}, \end{cases} \quad (14)$$

where $|\mathbf{u}| = n + m + k + l$ and $\mathbf{u} <' \mathbf{v}$ means that the first nonzero difference $r - k, s - l, p - n, q - m$ is positive. The resulting ordered sequence of the moments starts as follows:

$$1, \langle \hat{a} \rangle, \langle \hat{a}^\dagger \rangle, \langle \hat{b} \rangle, \langle \hat{b}^\dagger \rangle, \langle \hat{a}^2 \rangle, \langle \hat{a}^\dagger \hat{a} \rangle, \langle \hat{a}^{\dagger 2} \rangle, \langle \hat{a} \hat{b} \rangle, \langle \hat{a}^\dagger \hat{b} \rangle, \langle \hat{b}^2 \rangle, \langle \hat{a} \hat{b}^\dagger \rangle, \langle \hat{a}^\dagger \hat{b}^\dagger \rangle, \langle \hat{b}^\dagger \hat{b} \rangle, \langle \hat{b}^{\dagger 2} \rangle, \dots \quad (15)$$

One can easily check that the moments $M_{pqrs,nmkl}$, Eq. (13), can be expressed in terms of the moments of the original state as follows

$$\langle \hat{a}^{\dagger q} \hat{a}^p \hat{a}^{\dagger n} \hat{a}^m \hat{b}^{\dagger s} \hat{b}^r \hat{b}^{\dagger k} \hat{b}^l \rangle^{\text{PT}} = \langle \hat{a}^{\dagger q} \hat{a}^p \hat{a}^{\dagger n} \hat{a}^m \hat{b}^{\dagger l} \hat{b}^k \hat{b}^{\dagger r} \hat{b}^s \rangle.$$

Note that these moments are linear combinations of normally-ordered ones, $\langle \hat{a}^{\dagger n} \hat{a}^m \hat{b}^{\dagger k} \hat{b}^l \rangle$, according to the following formula for the antinormally-ordered product $\hat{a}^n \hat{a}^{\dagger m}$,

$$\hat{a}^n \hat{a}^{\dagger m} = \sum_{k=0}^{\min(n,m)} \frac{n!m!}{k!(n-k)!(m-k)!} \hat{a}^{\dagger m-k} \hat{a}^{n-k}. \quad (16)$$

The nonnegativity of all the minors of the quadratic form (12) can now be written as

$$\forall N : \quad D_N = \begin{vmatrix} M_{11} & M_{12} & \dots & M_{1N} \\ M_{21} & M_{22} & \dots & M_{2N} \\ \dots & \dots & \dots & \dots \\ M_{N1} & M_{N2} & \dots & M_{NN} \end{vmatrix} \geq 0, \quad (17)$$

where the moments M_{ij} defined in Eq. (13) are now given as

$$M_{ij} = \langle \hat{a}^{\dagger q} \hat{a}^p \hat{a}^{\dagger n} \hat{a}^m \hat{b}^{\dagger l} \hat{b}^k \hat{b}^{\dagger r} \hat{b}^s \rangle, \quad (18)$$

with (n, m, k, l) and (p, q, r, s) being the i th and the j th indices in the ordered sequence (15). Now we can explicitly formulate a necessary and sufficient condition for the positivity of partial transposition. *The partial transposition of a bipartite quantum state is nonnegative, if and only if all the determinants*

$$D_N = \begin{vmatrix} 1 & \langle \hat{a} \rangle & \langle \hat{a}^\dagger \rangle & \langle \hat{b}^\dagger \rangle & \langle \hat{b} \rangle & \dots \\ \langle \hat{a}^\dagger \rangle & \langle \hat{a}^\dagger \hat{a} \rangle & \langle \hat{a}^{\dagger 2} \rangle & \langle \hat{a}^\dagger \hat{b}^\dagger \rangle & \langle \hat{a}^\dagger \hat{b} \rangle & \dots \\ \langle \hat{a} \rangle & \langle \hat{a}^2 \rangle & \langle \hat{a} \hat{a}^\dagger \rangle & \langle \hat{a} \hat{b}^\dagger \rangle & \langle \hat{a} \hat{b} \rangle & \dots \\ \langle \hat{b} \rangle & \langle \hat{a} \hat{b} \rangle & \langle \hat{a}^\dagger \hat{b} \rangle & \langle \hat{b}^\dagger \hat{b} \rangle & \langle \hat{b}^2 \rangle & \dots \\ \langle \hat{b}^\dagger \rangle & \langle \hat{a} \hat{b}^\dagger \rangle & \langle \hat{a}^\dagger \hat{b}^\dagger \rangle & \langle \hat{b}^{\dagger 2} \rangle & \langle \hat{b} \hat{b}^\dagger \rangle & \dots \end{vmatrix} \quad (19)$$

are nonnegative, that is

$$\forall N : \quad D_N \geq 0. \quad (20)$$

The other way around, if there exists a negative determinant,

$$\exists N : \quad D_N < 0, \quad (21)$$

the NPT has been demonstrated. Hence the condition (21) is a necessary and sufficient condition for the NPT. That is, it provides a sufficient condition for the state under consideration to be entangled. This result is the central finding of the present contribution. It is important to stress that all the moments occurring in our conditions can be determined experimentally. This requires a straightforward extension of the method, as proposed for measuring the moments of a single-mode radiation field [11], to the case of two modes.

Even though the above conditions (21) are necessary and sufficient for NPT, for practical applications it may be useful to derive other types of conditions. They may be more efficient for verifying entanglement of particular quantum states in practice, violations of these new conditions may be found easier than the negativities of the determinants (19). For this purpose we consider the following two possibilities:

- (i) *Restricting the generality* of the operator (11) by restricting the generality of the coefficients c_{nmkl} leads to other conditions for inseparability. For example, setting some coefficients c_{nmkl} to zero leads to conditions for the nonnegativity of subdeterminants of D_N , which are obtained by canceling rows and columns with the same indices.
- (ii) *Identifying some coefficients* c_{nmkl} leads to another way to formulate new conditions for NPT by reducing the number of independent variables of the form (12). For example, the quadratic form derived

for the operator $\hat{f} = c_1 + c_2 \hat{a} + c_3 \hat{b}$ depends on three independent (complex) variables c_1 , c_2 and c_3 . If we identify the last two coefficients $c_2 = c_3$, i.e. $\hat{f} = c_1 + c_2(\hat{a} + \hat{b})$, we obtain a quadratic form of two variables c_1 and c_2 . The resulting sufficient conditions for NPT can be formulated in terms of the nonnegativity of determinants whose entries are combinations of the moments (18).

Examples for the usefulness of both types of new conditions will be considered in the following.

Let us compare our approach with some previously known results. We start with the separability condition found by Simon [4]. Define matrices A_1 , A_2 , C and J by the expressions

$$A_i = \begin{pmatrix} \langle (\Delta \hat{x}_i)^2 \rangle & \langle \{ \Delta \hat{x}_i, \Delta \hat{p}_i \} \rangle \\ \langle \{ \Delta \hat{x}_i, \Delta \hat{p}_i \} \rangle & \langle (\Delta \hat{p}_i)^2 \rangle \end{pmatrix}, \quad (22)$$

$$C = \begin{pmatrix} \langle \Delta \hat{x}_1 \Delta \hat{x}_2 \rangle & \langle \Delta \hat{x}_1 \Delta \hat{p}_2 \rangle \\ \langle \Delta \hat{p}_1 \Delta \hat{x}_2 \rangle & \langle \Delta \hat{p}_1 \Delta \hat{p}_2 \rangle \end{pmatrix},$$

and $J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. The Simon condition can be formulated as follows: for any separable quantum state the quantity S ,

$$S = \det A_1 \det A_2 + \left(\frac{1}{4} + \det C \right)^2 - \text{Tr} \left(A_1 J C J A_2 J C^T J \right) - \frac{1}{4} (\det A_1 + \det A_2) \geq 0, \quad (23)$$

is nonnegative. Since the position and momentum operators $\hat{x}_i$ and $\hat{p}_i$, $i = 1, 2$, are linear combinations of the creation and annihilation operators, the condition (23) can also be formulated in terms of the moments M_{ij} , Eq. (18). By using our concept the left hand side of the inequality (23) reads much simpler in terms of the moments M_{ij} . Simon's quantity S is nothing but our fifth order determinant D_5 :

$$S \equiv D_5. \quad (24)$$

Therefore the Simon condition (23) is exactly one condition of our hierarchy (20).

Another separability condition has been formulated in [5]. For any non-zero real parameter r one may define operators $\hat{u}$ and $\hat{v}$ via

$$\hat{u} = |r| \hat{x}_1 + r^{-1} \hat{x}_2, \quad \hat{v} = |r| \hat{p}_1 - r^{-1} \hat{p}_2. \quad (25)$$

Then, for any separable state the inequality

$$\langle (\Delta \hat{u})^2 \rangle + \langle (\Delta \hat{v})^2 \rangle - (r^2 + r^{-2}) \geq 0 \quad (26)$$

holds true for all $r \neq 0$. The condition (26) can be simplified as follows. We express the operators $\hat{u}$ and $\hat{v}$ as linear combinations of the creation and annihilation operators and substitute them into the inequality (26). Now its left

hand side can be minimized with respect to r . Then the inequality (26) is found to be equivalent to

$$\langle \Delta \hat{a}^\dagger \Delta \hat{a} \rangle \langle \Delta \hat{b}^\dagger \Delta \hat{b} \rangle \geq \text{Re}^2 \langle \Delta \hat{a} \Delta \hat{b} \rangle. \quad (27)$$

The inequality (27) is related to a subdeterminant of D_N , to the one resulting from the operator $\hat{f} = c_1 + c_2 \hat{a} + c_3 \hat{b}$:

$$d = \begin{vmatrix} 1 & \langle \hat{a} \rangle & \langle \hat{b} \rangle \\ \langle \hat{a}^\dagger \rangle & \langle \hat{a}^\dagger \hat{a} \rangle & \langle \hat{a}^\dagger \hat{b} \rangle \\ \langle \hat{b} \rangle & \langle \hat{a} \hat{b} \rangle & \langle \hat{b}^\dagger \hat{b} \rangle \end{vmatrix}. \quad (28)$$

The condition $d \geq 0$ explicitly reads as

$$\langle \Delta \hat{a}^\dagger \Delta \hat{a} \rangle \langle \Delta \hat{b}^\dagger \Delta \hat{b} \rangle \geq |\langle \Delta \hat{a} \Delta \hat{b} \rangle|^2, \quad (29)$$

from which it becomes clear that the condition (26) is a weaker form of the nonnegativity of our determinant (28).

There have been various other results in the literature. We can only briefly comment on some of them. A generalization of the inequality (26) was obtained in [8]. The main result is not explicitly based on the nonnegativity of partial transposition, but in fact it is a consequence of it. Using our theory we can show that the main inequality follows from Eq. (10) for the operator $\hat{f} = c_1 + c_2 \hat{A}_1 + c_3 \hat{B}_1 + c_4 \hat{A}_2 + c_5 \hat{B}_2$ (in the notation of work [8]), which is readily obtained by our approach (ii). Another inequality has been obtained in [6]. It differs from the condition (26) and its generalization [8] in that it gives a lower bound for the product of the variances of the operators (25) rather than for their sum. Though this inequality does not explicitly use NPT, in fact it is also a consequence of it. The condition provided in [6] easily follows from the nonnegativity of d , Eq. (28), together with the determinant corresponding to the operator $\hat{f} = c_1 + c_2(\hat{a} + \hat{b}) + c_3(\hat{a}^\dagger + \hat{b}^\dagger)$. Yet another condition has been recently obtained in [7]. It is easy to verify that this condition is a weaker form of the nonnegativity of the determinant corresponding to the operator $\hat{f} = c_1 + c_2 \hat{a} \hat{b} + c_3 \hat{a}^\dagger \hat{b}^\dagger$.

Let us illustrate our theory for the example of an entangled quantum composed of two coherent states,

$$|\psi_-\rangle = \mathcal{N}_-(\alpha, \beta) \left(|\alpha, \beta\rangle - |-\alpha, -\beta\rangle \right), \quad (30)$$

where the normalization is

$$\mathcal{N}_-(\alpha, \beta) = \left(2(1 - e^{-2(|\alpha|^2 + |\beta|^2)}) \right)^{-1/2}. \quad (31)$$

The violations of the conditions (23) and (26) fail to demonstrate the entanglement of this state. However, we may apply the following simple subdeterminant:

$$s = \begin{vmatrix} 1 & \langle \hat{b}^\dagger \rangle & \langle \hat{a} \hat{b}^\dagger \rangle \\ \langle \hat{b} \rangle & \langle \hat{b}^\dagger \hat{b} \rangle & \langle \hat{a} \hat{b}^\dagger \hat{b} \rangle \\ \langle \hat{a}^\dagger \hat{b} \rangle & \langle \hat{a}^\dagger \hat{b}^\dagger \hat{b} \rangle & \langle \hat{a}^\dagger \hat{a} \hat{b}^\dagger \hat{b} \rangle \end{vmatrix}. \quad (32)$$

Explicitly it reads as

$$s = -|\alpha|^2 |\beta|^4 \frac{\coth(|\alpha|^2 + |\beta|^2)}{\sinh^2(|\alpha|^2 + |\beta|^2)}, \quad (33)$$

which is clearly negative for all non-zero parameters and hence the entanglement is verified.

Finally, we will consider another special kind of separability condition obtained by our approach (i). Let us consider the simplest subdeterminants of the D_N , namely those of second order. They correspond to the condition (10) with $\hat{f}$ being a linear combination of two terms,

$$\hat{f} = c_1 \hat{a}^{\dagger n} \hat{a}^m \hat{b}^{\dagger k} \hat{b}^l + c_2 \hat{a}^{\dagger p} \hat{a}^q \hat{b}^{\dagger r} \hat{b}^s. \quad (34)$$

Explicitly the resulting separability condition reads as

$$\begin{aligned} & \langle \hat{a}^{\dagger m} \hat{a}^n \hat{a}^{\dagger n} \hat{a}^m \hat{b}^{\dagger l} \hat{b}^k \hat{b}^{\dagger k} \hat{b}^l \rangle \langle \hat{a}^{\dagger q} \hat{a}^p \hat{a}^{\dagger p} \hat{a}^q \hat{b}^{\dagger s} \hat{b}^r \hat{b}^{\dagger r} \hat{b}^s \rangle \geq \\ & \left| \langle \hat{a}^{\dagger m} \hat{a}^n \hat{a}^{\dagger p} \hat{a}^q \hat{b}^{\dagger s} \hat{b}^r \hat{b}^{\dagger k} \hat{b}^l \rangle \right|^2. \end{aligned} \quad (35)$$

This leads to rather simple conditions including moments of higher orders. Note that the conditions (35) generalize those derived recently in [12].

In conclusion, we have derived a hierarchy of conditions that is necessary and sufficient for the negativity of the partial transposition of the density operator of a bipartite harmonic oscillator. In fact this provides a sufficient condition for entanglement in terms of observable quantities, that can be used for a variety of practical applications. We have also demonstrated that several previously known conditions appear to be special cases of our approach.

* Electronic address: evgeny.shchukin@uni-rostock.de

† Electronic address: werner.vogel@uni-rostock.de

- [1] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996).
- [2] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Lett. A **223**, 1 (1996).
- [3] P. Horodecki, Phys. Lett. A **232**, 333 (1997).
- [4] R. Simon, Phys. Rev. Lett. **84**, 2726 (2000).
- [5] L.-M. Duan, G. Giedke, J. I. Cirac, and P. Zoller, Phys. Rev. Lett. **84**, 2722 (2000).
- [6] S. Mancini, V. Giovannetti, D. Vitali, and P. Tombesi, Phys. Rev. Lett. **88**, 120401 (2002).
- [7] G. S. Agarwal and A. Biswas, quant-ph/0501012 (2005).
- [8] M. G. Raymer, A. C. Funk, B. C. Sanders, and H. de Guise, Phys. Rev. A **67**, 052104 (2003).
- [9] T. Richter and W. Vogel, Phys. Rev. Lett. **89**, 283601 (2002).
- [10] E. Shchukin, T. Richter, and W. Vogel, Phys. Rev. A **71**, 011802(R) (2005).
- [11] E. Shchukin and W. Vogel, quant-ph/0506029 (2005), Phys. Rev. A, in press.
- [12] M. Hillery and M. Zubairy, quant-ph/0507168 (2005).